EPHOU 02-007

arXiv:gr-qc/0211056v3 24 Dec 2002

Analytic Evaluation of the Decay Rate for an Accelerated Proton

Hisao Suzuki* and Kunimasa Yamada†
Department of Physics, Hokkaido University, Sapporo 060-0810, Japan

(Dated: February 3, 2008)

We evaluate the decay rate of the uniformly accelerated proton. We obtain an analytic expression for inverse beta decay process caused by the acceleration. We evaluate the decay rate both from the inertial frame and from the accelerated frame where we should consider thermal radiation by Unruh effect. We explicitly check that the decay rates obtained in both frame coincide with each other.

PACS numbers: 13.30.-a, 04.62.+v, 12.15.Ji, 14.20.Dh

I. INTRODUCTION

In 1974 Hawking announced a celebrated result that the quantum thermal radiance occurs from black hole [1], which is the most interesting result of the quantum field theory in curved space-time. Since then, the study of the field theory has been of much interest. It appeared that the quantum radiation is not special phenomenon in strong gravitation but analogous effect exists even in the flat space-time [2]. The effect is caused by the fact that there is no vacuum which is invariant under the general coordinate transformations although the vacuum is invariant under the special coordinate transformations. A typical example is the accelerated observer which feels thermal radiation of the particles depending on the rate of acceleration. The effects in the accelerated system are not hard to be analyzed and is more tractable than the purely gravitational effects.

This phenomenon in flat space-time was shown by Fulling-Davies-Unruh in 1973, which is now called Unruh effect [3, 4]. A vacuum in rest frame is not a vacuum in accelerated systems. Actually, the vacuum in rest frames is a thermal bath in the accelerated frames [5]. This means that the definition of particle in quantum field theory depends on the observer and a strongly curved space-time by the acceleration corresponds to a strong gravitation.

Until now, many phenomena which the acceleration gives the interesting influence have been shown in the cosmological setting [6]. There is only one experiment about Unruh effect by Chen and Tajima [7]. They found that an electron which is accelerated by electronic field quivered. They showed that the random absorption of quanta from the FDU thermal bath cause this quivered. However, the experiment of decay is not existing.

More recently, the decay of accelerated particles was studied. In 1997 R. Muller [8] estimated the decay rates of the accelerated particle [9]. However, it was shown that decay rate is very small in our experimental setup.

It was G. E. A Matsas and D. A. T. Vanzella [10, 11] who realized that the decay rate can be used as a "*theoretical check*" of Unruh effect. The most interesting decay is in the case where we can deal with a particle which doesn't decay in inertial frame but can in accelerated frame. A typical example is for the proton. In inertial frame the accelerated proton can decay, and we can write the process as

$$\Gamma_{\rm rest}; \qquad p^+ \to n + e^+ + \nu_e .$$

In the accelerated frame, the observer takes the same acceleration as proton then the proton is stable but his space is in a thermal bath of particles. So the proton should absorb e^- and $\bar{\nu}$, and the processes become the inverse β decays as

$$\begin{aligned} \Gamma_{\rm acc}; \qquad & p^+ + e^- \to n + \nu, \\ & p^+ + \bar{\nu} \ \ \to n + e^+, \\ & p^+ + e^- + \bar{\nu} \to n. \end{aligned}$$

Both of the cross sections should agree with each other because they are the probabilities of the identical physical phenomenon and it is independ on the frames we use. However, the calculation is very complicating because the solutions of the Dirac equation in accelerated frame become a special function even if we represent protons by the classical current and restrict to tree level. Therefore, they restricted the system to two-dimensions. They could represent the decay rate analytically for Minkowski frame in the limit where neutrino is massless. However, the decay rate for the accelerated frame could not be evaluated analytically. Therefore, they showed the agreement numerically for massless neutrino. This calculation shows that the Unruh effect is inevitable for the accelerated frame. An interesting point of their approach is that they use a decay rate, true physical observables, for showing the existence of Unruh effect although the effect itself was established earlier (see Ref. [6] for Review).

They used two dimensional setting and treated neutrino for massless field for simplicity. However, it is desirable if we can show that decay rates calculated in both frame are identical even for massive neutrino in four dimensions because massiveness of neutrino is now proved experimentally.

The main aim of this paper is to show that we can complete their calculation by showing that the decay rate of

---

*Electronic address: hsuzuki@phys.sci.hokdai.ac.jp
†Electronic address: kunimasa@particle.sci.hokudai.ac.jp

proton is independent of the frame not by numerical computation but by analytical computation. Moreover, We will perform the evaluation in four dimensional setting and treat neutrino a massive field. It is interesting that the decay rate can be analyzed analytically.

In Section II we will calculate the cross section of β decay in the inertial frame. We will show that the decay rate can be obtained analytically in terms of a function which is an analog of Meijer's G-function of two variables.

In Section III, we will perform the calculation in accelerating frame. We will be able to check that the resulting function is identical to the one obtained in inertial frame.

Section IV is devoted to the discussions.

In Appendix B, we list the explicit form of the function appeared in our main result.

II. INERTIAL FRAME

In this section, we analysis the β decay of the accelerated proton in the inertial frame. In this frame, the accelerated proton decay resulting from the acceleration.

In the flat space-time, the solutions of the Dirac equation are simple but the calculation becomes complicate because of the vector current is on the hyperbola. We will mainly follow the notation appeared in Ref. [10, 11]

A. Accelerated Proton current

There may be several methods to represent the acceleration. One way is the followings. We represent the proton as a classical current. The position of the current should be on the world line of the accelerated particle, "hyperbola". To do this, we introduce the Rindler coordinates. Rindler coordinates correspond to the world line of the uniformly accelerated observers. The inertial coordinates are shown by (x^0, x^1, x^2, x^3) and the Rindler coordinates which show acceleration along the 3-axis are (v, x^1, x^2, u) with $0 < u < \infty$ and $-\infty < v < \infty$. Two coordinates are related by

$$\begin{aligned} x^0 &= u \sinh v, \\ x^1 &= x^1, \\ x^2 &= x^2, \\ x^3 &= u \cosh v. \end{aligned} \tag{1}$$

Then the line element is described in the Rindler coordinates by

$$ds^2 = u^2 dv^2 - (dx^1)^2 - (dx^2)^2 - du^2. \tag{2}$$

If some particle uniformly accelerated with a proper acceleration a, then $u = a^{-1} = \text{const.}$ is its world line. As a result, the protons which are accelerated to the axis of x^3 on $x^1 = 0,\ x^2 = 0$ are represented as following classical current

$$j^\mu = q u^\mu \delta(x^1)\delta(x^2)\delta(u - a^{-1}), \tag{3}$$

where q is a small coupling constant and u^μ is the four-velocity $u^\mu = (a, 0, 0, 0)$ in Rindler coordinates. To deal with the proton-decay, nucleons $|n\rangle$ and protons $|p\rangle$ are represented as excited and unexcited states of the nucleon, respectively. For Hamiltonian $\hat{H}$, neutron and proton mass m_n and m_p, respectively, we set

$$\hat{H}|n\rangle = m_n|n\rangle, \quad \hat{H}|p\rangle = m_p|p\rangle. \tag{4}$$

We replace q in last current by the Hermitian monopole

$$\hat{q}(\tau) \equiv e^{i\hat{H}\tau}\hat{q}(0)e^{-i\hat{H}\tau}, \tag{5}$$

where $\tau = v/a$ is proper time of the proton.

Then the current four-vector is

$$\hat{j}^\mu = \hat{q}(\tau) u^\mu \delta(x^1)\delta(x^2)\delta(u - a^{-1}). \tag{6}$$

This is clear that the particle is on the hyperbola.

B. Fermionic field quantization

For electrons and neutrinos, we write the fermionic fields satisfying the Dirac equation $(i\gamma^\mu\partial_\mu - m)\psi^{(\pm\omega)}_{\mathbf{k}\sigma} = 0$ as

$$\hat{\Psi}(x) = \sum_{\sigma=\pm}\int_{-\infty}^{\infty} d^3k \left[\hat{b}_{\mathbf{k}\sigma}\psi^{(+\omega)}_{\mathbf{k}\sigma}(x) + \hat{d}^\dagger_{\mathbf{k}\sigma}\psi^{(-\omega)}_{\mathbf{k}-\sigma}(x)\right], \tag{7}$$

where $x = (x^0, x^1, x^2, x^3)$, $k = (\omega, k^1, k^2, k^3)$, $\mathbf{x} = (x^1, x^2, x^3)$, $\mathbf{k} = (k^1, k^2, k^3)$. $\hat{b}_{\mathbf{k}\sigma}$ and $\hat{d}^\dagger_{\mathbf{k}\sigma}$ are annihilation and creation operators of fermions and antifermions, respectively. $\mathbf{k}$ and σ are momentum and polarization, respectively. $\psi^{(+\omega)}_{\mathbf{k}\sigma}$ and $\psi^{(-\omega)}_{\mathbf{k}\sigma}$ are positive and negative frequency solutions of the Dirac equation.

By solving the Dirac equation we obtain the orthonormal mode solutions [12],

$$\psi^{(\pm\omega)}_{\mathbf{k}\sigma}(x) = \frac{e^{\mp i k_\mu x^\mu}}{(2\pi)^{\frac{3}{2}}} u^{(\pm\omega)}_\sigma(\mathbf{k}), \tag{8}$$

where

$$u^{(\pm\omega)}_\sigma(\mathbf{k}) = \frac{k_\mu\gamma^\mu \pm m}{\sqrt{\omega(\omega \pm m)}}\hat{u}_\sigma \tag{9}$$

and

$$\hat{u}_+ = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \hat{u}_- = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}. \tag{10}$$

We choose the traditional inner product of the form

$$\begin{aligned} \langle\psi^{(\pm\omega)}_{\mathbf{k}\sigma}(x), \psi^{(\pm\omega')}_{\mathbf{k}'\sigma'}(x)\rangle &\equiv \int_\Sigma d\Sigma_\mu \psi^{(\pm\omega)}_{\mathbf{k}\sigma}(x)\gamma^\mu\psi^{(\pm\omega')}_{\mathbf{k}'\sigma'}(x) \\ &= \delta^3(\mathbf{k} - \mathbf{k}')\delta_{\sigma\sigma'}\delta_{\pm\omega\pm\omega'} \end{aligned} \tag{11}$$

and the solutions are normalized by this definition, where $\bar{\psi} \equiv \psi^\dagger \gamma^0$, $d\Sigma_\mu \equiv n_\mu d\Sigma$, n^μ is a unit vector orthogonal to Σ and we have chosen Σ to be the hypersurface of constant x^0. The creation and annihilation operators satisfy the equal-time anti-commutation relations:

$$\begin{aligned} \left\{\hat{b}_{\mathbf{k}\sigma}, \hat{b}^\dagger_{\mathbf{k}'\sigma'}\right\} &= \left\{\hat{d}_{\mathbf{k}\sigma}, \hat{d}^\dagger_{\mathbf{k}'\sigma'}\right\} = \delta^3(\mathbf{k}-\mathbf{k}')\delta_{\sigma\sigma'}, \\ \left\{\hat{b}_{\mathbf{k}\sigma}, \hat{b}_{\mathbf{k}'\sigma'}\right\} &= \left\{\hat{d}_{\mathbf{k}\sigma}, \hat{d}_{\mathbf{k}'\sigma'}\right\} \\ &= \left\{\hat{b}_{\mathbf{k}\sigma}, \hat{d}_{\mathbf{k}'\sigma'}\right\} = \left\{\hat{b}_{\mathbf{k}\sigma}, \hat{d}^\dagger_{\mathbf{k}'\sigma'}\right\} = 0. \end{aligned} \tag{12}$$

Using electron $\hat{\Psi}_e$ and neutrino $\hat{\Psi}_\nu$ fields, we write the Fermi action by the nucleon current (2.6)

$$\hat{S}_I = \int d^4x\sqrt{-g}\hat{j}_\mu(\hat{\bar{\Psi}}_\nu\gamma^\mu\hat{\Psi}_e + \hat{\bar{\Psi}}_e\gamma^\mu\hat{\Psi}_\nu), \tag{13}$$

where the first term is used by inverse β decay.

You can see these general formation on any book of Quantum Field Theory in the inertial frame. We are ready to start to calculate the cross section of the accelerated proton in the inertial frame.

C. Calculation of Cross Section

We are now going to calculate the cross section of the β decay using the current and field in the subsection A. The way of deriving the cross section is normal.

First, the vacuum transition amplitude of the proton decay is written by

$$\mathcal{A}^{p\to n} = \langle n| \otimes \langle e^+_{k_e\sigma_e}, \nu_{k_\nu\sigma_\nu}|\hat{S}_I|0\rangle \otimes |p\rangle. \tag{14}$$

It is straightforward to compute $\mathcal{A}^{p\to n}$ for the Fermi action $\hat{S}_I$ and we obtain

$$\begin{aligned} \mathcal{A}^{p\to n} &= G_F \int_{-\infty}^{\infty} d(x^0)d(x^3)\frac{e^{i\Delta m\tau}u_\mu}{\sqrt{a^2(x^0)^2+1}} \\ &\times\delta(x^3-\sqrt{(x^0)^2+a^{-2}})\langle e^+_{k_e\sigma_e}, \nu_{k_\nu\sigma_\nu}|\hat{\bar{\Psi}}_\nu\gamma^\mu\hat{\Psi}_e|0\rangle, \end{aligned} \tag{15}$$

where $\Delta m \equiv m_n - m_p$, $\tau = a^{-1}\sinh^{-1}(ax^0)$ is the nucleon's proper time, $G_F \equiv \langle p|\hat{q}(0)|n\rangle$ is the Fermi constant. We can substitute the field and integrate by x^3. The differential transition rate is

$$\frac{d^6\mathcal{P}^{p\to n}_{\text{in}}}{d^3k_e d^3k_\nu} = \sum_{\sigma_e}\sum_{\sigma_\nu}|\mathcal{A}^{p\to n}|^2\,. \tag{16}$$

We obtain it by integration of proper times τ_1 and τ_2

$$\begin{aligned} \frac{d^6\mathcal{P}^{p\to n}_{\text{in}}}{d^3k_e d^3k_\nu} &= \frac{G_F^2}{(2\pi)^6}\int_{-\infty}^{\infty} d\tau_1 d\tau_2\, u_\mu u^*_\nu \sum_{\sigma_e}\sum_{\sigma_\nu}\left[\bar{u}^{(+\omega_\nu)}_{\sigma_\nu}\gamma^\mu u^{(-\omega_e)}_{-\sigma_e}\right]\left[\bar{u}^{(+\omega_\nu)}_{\sigma_\nu}\gamma^\nu u^{(-\omega_e)}_{-\sigma_e}\right]^* \\ &\times\exp i\left[\Delta m(\tau_1-\tau_2) + a^{-1}(\omega_\nu+\omega_e)(\sinh a\tau_1 - \sinh a\tau_2) - a^{-1}(k^3_\nu + k^3_e)(\cosh a\tau_1 - \cosh a\tau_2)\right]. \end{aligned} \tag{17}$$

Now in order to calculate the spin sums, we use the following standard formula:

$$\begin{aligned} &\sum_{\sigma_\alpha}\sum_{\sigma_\beta}\left[\bar{u}^{(\pm\omega)}_{\sigma_\alpha}\Gamma_1 u^{(\pm\omega)}_{\sigma_\beta}\right]\left[\bar{u}^{(\pm\omega)}_{\sigma_\alpha}\Gamma_2 u^{(\pm\omega)}_{\sigma_\beta}\right]^* \\ &= \text{Tr}\left[\Gamma_1\sum_{\sigma_\alpha}u^{(\pm\omega)}_{\sigma_\alpha}\bar{u}^{(\pm\omega)}_{\sigma_\alpha}\gamma^0\Gamma_2^\dagger\gamma^0\sum_{\sigma_\beta}u^{(\pm\omega)}_{\sigma_\beta}\bar{u}^{(\pm\omega)}_{\sigma_\beta}\right], \end{aligned} \tag{18}$$

where α and β represent e or ν.

The completeness relations can be written as

$$\sum_{\sigma_\alpha}u^{(\pm\omega_\alpha)}_{\sigma_\alpha}(\mathbf{k}_\alpha)\bar{u}^{(\pm\omega_\alpha)}_{\sigma_\alpha}(\mathbf{k}_\alpha) = \frac{1}{2\omega_\alpha}(k_{\alpha\mu}\gamma^\mu \pm m_\alpha) \tag{19}$$

and we introduce s and ξ by

$$\tau_1 = s + \frac{\xi}{2},\;\; \tau_2 = s - \frac{\xi}{2}. \tag{20}$$

By using the spin sum

$$\begin{aligned} &\sum_{\sigma_e}\sum_{\sigma_\nu}\left[\bar{u}^{(+\omega_\nu)}_{\sigma_\nu}\gamma^\mu u^{(-\omega_e)}_{-\sigma_e}\right]\left[\bar{u}^{(+\omega_\nu)}_{\sigma_\nu}\gamma^\nu u^{(-\omega_e)}_{-\sigma_e}\right]^* \\ &= \frac{1}{\omega_e\omega_\nu}\left[(\omega_e\omega_\nu + k^3_e k^3_\nu)\cosh 2as \right. \\ &\quad \left. -(\omega_e k^3_\nu + k^3_e\omega_\nu)\sinh 2as - m_e m_\nu\cosh a\xi\right] \\ &\qquad +(\text{odd function of } k^1_\alpha \text{ and } k^2_\alpha) \end{aligned} \tag{21}$$

and the change of variables

$$k'^1_\alpha = k^1_\alpha,\; k'^2_\alpha = k^2_\alpha,\; k'^3_\alpha = -\omega_\alpha\sinh as + k^3_\alpha\cosh as, \tag{22}$$

we can obtain the differential transition rate

$$\frac{1}{T}\frac{d^6\mathcal{P}^{p\to n}_{\rm in}}{d^3k'_e d^3k'_\nu} = \frac{G_F^2}{(2\pi)^6}\frac{1}{\omega'_e\omega'_\nu}\int_{-\infty}^{\infty} d\xi e^{i[\Delta m\xi + \frac{\omega'_e+\omega'_\nu}{2a}\sinh\frac{a\xi}{2}]} \times(\omega'_e\omega'_\nu + {k'}^3_e{k'}^3_\nu - m_e m_\nu\cosh a\xi), \tag{23}$$

where $T \equiv \int_{-\infty}^{\infty} ds$. To integrate by ξ we redefine new variable by

$$\lambda \equiv e^{\frac{a\xi}{2}}. \tag{24}$$

And we use the following notations

$$\widetilde{k}_\alpha \equiv \frac{k'_\alpha}{a},\ \ \widetilde{\omega}_\alpha \equiv \frac{\omega_\alpha}{a},\ \ \widetilde{m}_\alpha \equiv \frac{m_\alpha}{a},\ \ \widetilde{\Delta m} \equiv \frac{\Delta m}{a}. \tag{25}$$

Then the cross section is given in the form:

$$\begin{aligned}\Gamma^{p\to n}_{\rm in} &= \frac{1}{T}\int d\mathcal{P}^{p\to n}_{\rm in} \\ &= \frac{a^5G_F^2}{2^5\pi^6}\int_{-\infty}^{\infty}\frac{d^3\widetilde{k}_e d^3\widetilde{k}_\nu}{\widetilde{\omega}_e\widetilde{\omega}_\nu}d\lambda\, e^{i(\widetilde{\omega}_e+\widetilde{\omega}_\nu)(\lambda-\frac{1}{\lambda})} \\ &\quad\times\lambda^{2i\widetilde{\Delta m}-1}\left[\widetilde{\omega}_e\widetilde{\omega}_\nu - \frac{1}{2}\widetilde{m}_e\widetilde{m}_\nu(\lambda^2+\frac{1}{\lambda^2})\right].\end{aligned} \tag{26}$$

The integral of λ can be readily expressed as modified Bessel to find

$$\begin{aligned}\Gamma^{p\to n}_{\rm in} = \frac{2^2a^5G_F^2}{\pi^6e^{\pi\widetilde{\Delta m}}}\int_0^\infty d^3\widetilde{k}_e d^3\widetilde{k}_\nu\Bigg[K_{2i\widetilde{\Delta m}}\Big(\,2(\widetilde{\omega}_e+\widetilde{\omega}_\nu)\Big) \\ +\frac{1}{2}\frac{\widetilde{m}_e\widetilde{m}_\nu}{\widetilde{\omega}_e\widetilde{\omega}_\nu}\Big\{K_{2i\widetilde{\Delta m}+2}\Big(\,2(\widetilde{\omega}_e+\widetilde{\omega}_\nu)\Big) \\ +K_{2i\widetilde{\Delta m}-2}\Big(\,2(\widetilde{\omega}_e+\widetilde{\omega}_\nu)\Big)\Big\}\Bigg].\end{aligned} \tag{27}$$

However, this form is difficult to integrate by $\widetilde{k}_e$ and $\widetilde{k}_\nu$. Therefore, we use the integration formula of modified Bessel

$$K_\mu(z) = \frac{1}{2}\int_{C_1}\frac{ds}{2\pi i}\Gamma(-s)\Gamma(-s-\mu)\left(\frac{z}{2}\right)^{2s+\mu}. \tag{28}$$

It is not hard to show this formula in complex s plane by picking the residues of $\Gamma(-s)$ and $\Gamma(-s-\mu)$(see Appendix A). To evaluate Eq. (27) we use this formals and after some simple shift of variable we have

$$\begin{aligned}\Gamma^{p\to n}_{\rm in} = \frac{2a^5G_F^2}{\pi^6e^{\pi\widetilde{\Delta m}}}\int_0^\infty d^3\widetilde{k}_e d^3\widetilde{k}_\nu\int_{C_1}\frac{ds}{2\pi i}\left(\sqrt{\tilde{k}_e^2+\tilde{m}_e^2}+\sqrt{\tilde{k}_\nu^2+\tilde{m}_\nu^2}\right)^{2s+2i\widetilde{\Delta m}} \\ \times\left[\Gamma(-s)\Gamma(-s-2i\widetilde{\Delta m}) + \frac{1}{2}\frac{\widetilde{m}_e\widetilde{m}_\nu}{\widetilde{\omega}_e\widetilde{\omega}_\nu}\Big\{\Gamma(-s+1)\Gamma(-s-2i\widetilde{\Delta m}-1)+\Gamma(-s-1)\Gamma(-s-2i\widetilde{\Delta m}+1)\Big\}\right],\end{aligned} \tag{29}$$

where the contour C_1 must be chose so that all poles of $\Gamma(-s)\Gamma(-s-\mu)$ are picked up and must be selected as k integration don't infinity. But, this form of integration is still not simple for $\widetilde{k}_e$ and $\widetilde{k}_\nu$.

In order to perform the integration, we use the following expansion formula

$$(A+B)^z = \int_{C_2}\frac{dt}{2\pi i}\frac{\Gamma(-t)\Gamma(t-z)}{\Gamma(-z)}A^{t+z}B^t \tag{30}$$

to integrate easily by $\widetilde{k}_e$ and $\widetilde{k}_\nu$, where the contour C_2 is the path separating the poles of $\Gamma(-t)$ from those of $\Gamma(t-z)$ (see Appendix A).

Now we can rewrite Eq. (29) as

$$\begin{aligned}\Gamma^{p\to n}_{\rm in} &= \frac{a^5G_F^2}{2^5\pi^{\frac{7}{2}}e^{\pi\widetilde{\Delta m}}}\int_{C_s}\frac{ds}{2\pi i}\int_{C_t}\frac{dt}{2\pi i}\frac{(\tilde{m}_e^2)^s(\tilde{m}_\nu^2)^t}{\Gamma(-s-t+3)\Gamma(-s-t+\frac{7}{2})} \\ &\quad\times\Bigg[\Big|\Gamma(-s-t+i\widetilde{\Delta m}+3)\Big|^2\Gamma(-s)\Gamma(-t)\Gamma(-s+2)\Gamma(-t+2) \\ &\quad+{\rm Re}\Big\{\Gamma(-s-t+i\widetilde{\Delta m}+2)\Gamma(-s-t-i\widetilde{\Delta m}+4)\Big\}\Gamma(-s+\frac{1}{2})\Gamma(-t+\frac{1}{2})\Gamma(-s+\frac{3}{2})\Gamma(-t+\frac{3}{2})\Bigg]\end{aligned} \tag{31}$$

where the contour C_s and C_t is the path which picks up all poles of Gamma functions in s and t complex planes, respectively.

This is the two-dimensional analog of Meijer's G-function [13]. The explicit form can be obtained by evaluating contour integral. We list the results in the Appendix B.

III. ACCELERATED FRAME

In this section, we are going to analyze the same physical phenomenon view of accelerated system.

A. Fermionic field quantization

The Dirac equation in curved space-time is written by

$$(i\gamma_R^\mu \widetilde{\nabla}_\mu - m)\psi_{\mathbf{w}\sigma}^{(\omega)}(x) = 0, \tag{32}$$

where

$$\begin{aligned}
&x = (v, x^1, x^2, u),\\
&\mathbf{w} = (\omega, k^1, k^2), \;\; \mathbf{x} = (x^1, x^2, u),\\
&(e_0)^\mu = u^{-1}\delta_0^\mu, \; (e_i)^\mu = \delta_i^\mu,\\
&\gamma_R^\mu \equiv (e_\nu)^\mu \gamma^\nu,\\
&\widetilde{\nabla}_\mu \equiv \partial_\mu + \frac{1}{8}[\gamma^\alpha, \gamma^\beta](e_\alpha)^\lambda \nabla_\mu (e_\beta)_\lambda.
\end{aligned} \tag{33}$$

In the Rindler coordinates, the equation becomes

$$i\frac{\partial \psi_{\mathbf{w}\sigma}^{(\omega)}(x)}{\partial v} = \left(\gamma^0 m u - \frac{i\alpha^3}{2} - iu\alpha^i\partial_i\right)\psi_{\mathbf{w}\sigma}^{(\omega)}(x), \tag{34}$$

where $\alpha^i \equiv \gamma^0\gamma^i$.

The fermionic field can be expanded as

$$\hat{\Psi}(x) = \sum_{\sigma=\pm}\int_0^\infty d\omega \int_{-\infty}^\infty d^2k \left[\hat{b}_{\mathbf{w}\sigma}\psi_{\mathbf{w}\sigma}^{(+\omega)} + \hat{d}_{\mathbf{w}\sigma}^\dagger \psi_{\mathbf{w}-\sigma}^{(-\omega)}\right]. \tag{35}$$

We will the solution in the form

$$\psi_{\mathbf{w}\sigma}^{(\omega)} = \bar{f}_{\mathbf{w}\sigma}(\mathbf{x})e^{-i\omega v/a} \tag{36}$$

for $-\infty < \omega < \infty$.

The function $\bar{f}_{\mathbf{w}\sigma}$ is eigenstate of Hamiltonian as

$$\hat{H}\bar{f}_{\mathbf{w}\sigma} = \omega \bar{f}_{\mathbf{w}\sigma}, \tag{37}$$

where

$$\hat{H} = a\left[mu\gamma^0 - \frac{i\alpha^3}{2} - iu\alpha^i\partial_i\right]. \tag{38}$$

We denote two-component spinors χ_a by

$$\bar{f}_{\mathbf{w}\sigma}(\mathbf{x}) \equiv \begin{bmatrix} \bar{\chi}_1(\mathbf{x}, \mathbf{w}) \\ \bar{\chi}_2(\mathbf{x}, \mathbf{w}) \end{bmatrix}. \tag{39}$$

then we find that the Dirac equation takes the form

$$\delta^{ij}u\partial_i(u\partial_j\chi_1) = \left[m^2u^2 + \frac{1}{4} - \tilde{\omega}^2\right]\chi_1 - i\tilde{\omega}\sigma^3\chi_2, \tag{40a}$$

$$\delta^{ij}u\partial_i(u\partial_j\chi_2) = \left[m^2u^2 + \frac{1}{4} - \tilde{\omega}^2\right]\chi_2 - i\tilde{\omega}\sigma^3\chi_1, \tag{40b}$$

where i and j run over 1, 2, 3. By now, for squared equations we used $f_{\mathbf{w}\sigma}$, χ_1 and χ_2.

To simplify these equations, we introduce the functions $\phi^\pm \equiv \chi_1 \mp \chi_2$ and we can define $\xi^\pm$ and $\zeta^\pm$ through

$$\phi^\pm(\mathbf{x}, \mathbf{w}) \equiv \begin{bmatrix} \xi^\pm(\mathbf{x}, \mathbf{w}) \\ \zeta^\pm(\mathbf{x}, \mathbf{w}) \end{bmatrix}. \tag{41}$$

Given this definition, we can separate the equation for ξ and ζ in the form:

$$\delta^{ij}u\partial_i(u\partial_j\xi^\pm) = \left[m^2u^2 + \left(i\tilde{\omega} \pm \frac{1}{2}\right)^2\right]\xi^\pm, \tag{42a}$$

$$\delta^{ij}u\partial_i(u\partial_j\zeta^\pm) = \left[m^2u^2 + \left(i\tilde{\omega} \mp \frac{1}{2}\right)^2\right]\zeta^\pm. \tag{42b}$$

Here, we introduce ℓ as $\ell^2 = (k^1)^2 + (k^2)^2 + m^2$ by which the equation can be written as

$$\begin{aligned}
[\partial_1^2 + \partial_2^2 - m^2]\xi^\pm &= \frac{1}{u^2}\left[-u\partial_3(u\partial_3) + \left(i\tilde{\omega} \pm \frac{1}{2}\right)^2\right]\xi^\pm \\
&\equiv -\ell^2\xi^\pm.
\end{aligned} \tag{43}$$

The solution can be written in the form

$$\xi^\pm = A^\pm(\mathbf{w})\mathcal{X}(x^1, x^2, k^1, k^2)\mathcal{M}_{\mathbf{w}}^\pm(u). \tag{44}$$

This leads to the following decoupled equations

$$[\partial_1^2 + \partial_2^2 - m^2]\mathcal{X}(x^1, x^2, k^1, k^2) = 0, \tag{45}$$

$$\left[\frac{\partial^2}{\partial(\ell u)^2} + \frac{1}{\ell u}\frac{\partial}{\partial(\ell u)} - 1 - \left(\frac{i\tilde{\omega} \mp \frac{1}{2}}{\ell u}\right)^2\right]\mathcal{M}_{\mathbf{w}}^\pm(u) = 0. \tag{46}$$

We can readily solve the equation in the form

$$\mathcal{X}(x^1, x^2, k^1, k^2) = e^{ik_a x^a}, \tag{47}$$

$$\mathcal{M}_{\mathbf{w}}^\pm(u) = K_{i\tilde{\omega}\mp\frac{1}{2}}(\ell u), \tag{48}$$

where the index a run over 1 and 2.

Using the arbitrary function $A^\pm$ and $B^\pm$ of k, we find the most general solution can be written as

$$\xi^\pm = A^\pm(\mathbf{w})e^{ik_a x^a}K_{i\tilde{\omega}\mp\frac{1}{2}}(\ell u), \tag{49a}$$

$$\zeta^\pm = B^\pm(\mathbf{w})e^{ik_a x^a}K_{i\tilde{\omega}\pm\frac{1}{2}}(\ell u), \tag{49b}$$

which implies

$$f_{\mathbf{w}\sigma} = \frac{1}{2}\begin{bmatrix} \xi^+ + \xi^- \\ \zeta^+ + \zeta^- \\ -\xi^+ + \xi^- \\ -\zeta^+ + \zeta^- \end{bmatrix} = \frac{e^{ik_a x^a}}{2}\begin{bmatrix} A^+ K_{i\tilde{\omega}-\frac{1}{2}}(\ell u) + A^- K_{i\tilde{\omega}+\frac{1}{2}}(\ell u) \\ B^+ K_{i\tilde{\omega}+\frac{1}{2}}(\ell u) + B^- K_{i\tilde{\omega}-\frac{1}{2}}(\ell u) \\ -A^+ K_{i\tilde{\omega}-\frac{1}{2}}(\ell u) + A^- K_{i\tilde{\omega}+\frac{1}{2}}(\ell u) \\ -B^+ K_{i\tilde{\omega}+\frac{1}{2}}(\ell u) + B^- K_{i\tilde{\omega}-\frac{1}{2}}(\ell u) \end{bmatrix}. \tag{50}$$

It is easy to see that $f_{\mathbf{w}\sigma}$ satisfies

$$\left[i\alpha^i\partial_i - m\gamma^0 + \left(\frac{i\alpha^3}{2} + \tilde{\omega}\right)\frac{1}{u}\right] \times \left[i\alpha^j\partial_j - m\gamma^0 + \left(\frac{i\alpha^3}{2} - \tilde{\omega}\right)\frac{1}{u}\right] f_{\mathbf{w}\sigma} = 0. \tag{51}$$

Therefore, the solutions of Dirac equation are simply given by

$$\bar{f}_{\mathbf{w}\sigma} \equiv \left[i\alpha^i\partial_i - m\gamma^0 + \left(\frac{i\alpha^3}{2} - \tilde{\omega}\right)\frac{1}{u}\right] f_{\mathbf{w}\sigma}. \tag{52}$$

If we set

$$A^+ = B^- = 0 \tag{53}$$

we can find a normalized A^- and B^+ by letting

$$A^- = B^+ = \left[\frac{\cosh\tilde{\omega}\pi}{\pi\tilde{\ell}}\right]^{\frac{1}{2}} \equiv N, \tag{54}$$

where

$$\tilde{\ell} \equiv \frac{\ell}{a}, \tag{55}$$

which is normalized with respect to the inner product

$$\langle\psi^{(\omega)}_{\mathbf{w}\sigma}(x), \psi^{(\omega')}_{\mathbf{w}'\sigma'}(x)\rangle \equiv \int_\Sigma d\Sigma_\mu \psi^{(\omega)}_{\mathbf{w}\sigma}(x)\gamma^\mu_R \psi^{(\omega')}_{\mathbf{w}'\sigma'}(x) = \delta^3(\mathbf{w}-\mathbf{w}')\delta_{\sigma\sigma'}, \tag{56}$$

where $\bar{\psi} \equiv \psi^\dagger\gamma^0$ and Σ is set to be $v =$ const.. In this way, we find concrete form of fermionic field is written by

$$\psi^{(\omega)}_{\mathbf{w}\sigma}(x) = \frac{e^{-i\omega v/a + ik_a x^a}}{(2\pi)^{\frac{3}{2}}} u^{(\omega)}_\sigma(u, \mathbf{w}), \tag{57}$$

where

$$u^{(\omega)}_\sigma(u,\mathbf{w}) = N\gamma^0\left[(\tilde{k}_a\gamma^a + \tilde{m})K_{i\tilde{\omega}+\frac{1}{2}}(\ell u) + i\tilde{\ell}\gamma^3 K_{i\tilde{\omega}-\frac{1}{2}}(\ell u)\right]\hat{u}_\sigma \tag{58}$$

and

$$\hat{u}_+ = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \quad \hat{u}_- = \begin{bmatrix} 0 \\ 1 \\ 0 \\ -1 \end{bmatrix}. \tag{59}$$

The creation and annihilation operators should obey

$$\begin{aligned} \left\{\hat{b}_{\mathbf{w}\sigma}, \hat{b}^\dagger_{\mathbf{w}'\sigma'}\right\} &= \left\{\hat{d}_{\mathbf{w}\sigma}, \hat{d}^\dagger_{\mathbf{w}'\sigma'}\right\} = \delta^3(\mathbf{w}-\mathbf{w}')\delta_{\sigma\sigma'}, \\ \left\{\hat{b}_{\mathbf{w}\sigma}, \hat{b}_{\mathbf{w}'\sigma'}\right\} &= \left\{\hat{d}_{\mathbf{w}\sigma}, \hat{d}_{\mathbf{w}'\sigma'}\right\} \\ &= \left\{\hat{b}_{\mathbf{w}\sigma}, \hat{d}_{\mathbf{w}'\sigma'}\right\} = \left\{\hat{b}_{\mathbf{w}\sigma}, \hat{d}^\dagger_{\mathbf{w}'\sigma'}\right\} = 0. \end{aligned} \tag{60}$$

B. Calculation of Cross Section

The process of β decay in the accelerated frame looks very different from that in rest frame. In this case, a proton is stable but the whole space is FDU thermal bath characterized by a temperature $T = a/2\pi$. Therefore, proton absorbs e^- and $\bar{\nu}$ from FDU thermal bath and do emit e^+ and ν. Three processes are possible through the precesses:

(i) $\mathrm{p}^+ + \mathrm{e}^- \rightarrow \mathrm{n} + \nu$,
(ii) $\mathrm{p}^+ + \bar{\nu} \rightarrow \mathrm{n} + \mathrm{e}^+$,
(iii) $\mathrm{p}^+ + \mathrm{e}^- + \bar{\nu} \rightarrow \mathrm{n}$.

The transition rate is a combination of them.

Formally, we can calculate the cross sections by same way of the inertial system but we have to deal with three processes.

The transition amplitudes are written by

$$\begin{aligned} \mathcal{A}^{p\rightarrow n}_{\text{(i)}} &= \langle n| \otimes \langle\nu_{k_\nu\sigma_\nu}|\hat{S}_I|e^-_{k_{e^-}\sigma_{e^-}}\rangle \otimes |p\rangle \\ &= \frac{G_F}{a}\int_{-\infty}^{\infty} dv e^{i\Delta m v}\langle\nu_{k_\nu\sigma_\nu}|\hat{\bar{\Psi}}_\nu\gamma^0\hat{\Psi}_e|e^-_{k_{e^-}\sigma_{e^-}}\rangle \\ &= \frac{G_F}{2\pi}\delta(\omega_{e^-} - \omega_\nu - \Delta m)\bar{u}^{(\omega_\nu)}_{\sigma_\nu}\gamma^0 u^{(\omega_{e^-})}_{\sigma_{e^-}}, \end{aligned} \tag{61a}$$

$$\mathcal{A}^{p\rightarrow n}_{\text{(ii)}} = \frac{G_F}{2\pi}\delta(\omega_{\bar{\nu}} - \omega_{e^+} - \Delta m)\bar{u}^{(\omega_{e^+})}_{\sigma_{e^+}}\gamma^0 u^{(\omega_{\bar{\nu}})}_{\sigma_{\bar{\nu}}}, \tag{61b}$$

$$\mathcal{A}^{p\rightarrow n}_{\text{(iii)}} = \frac{G_F}{2\pi}\delta(\omega_{e^-} + \omega_{\bar{\nu}} - \Delta m)\bar{u}^{(-\omega_{\bar{\nu}})}_{-\sigma_{\bar{\nu}}}\gamma^0 u^{(\omega_{e^-})}_{\sigma_{e^-}}, \tag{61c}$$

where $\hat{S}_I$ is replaced γ^μ in inertial frame with γ^μ_R and $u^\mu = (a, 0, 0, 0)$.

We assume that the observer is in the thermal bath. We attach the fermionic thermal factor for each process. Then the differential transition rate per absorbed and emitted particle energies for each processes are written as

$$\begin{aligned}
&\frac{1}{T}\frac{d^6\mathcal{P}^{p\to n}_{\text{(i)}}}{d\omega_{e^-}d\omega_\nu d^2k_{e^-}d^2k_\nu} \\
&= \frac{1}{T}\sum_{\sigma_{e^-}}\sum_{\sigma_\nu}\left|\mathcal{A}^{p\to n}_{\text{(i)}}\right|^2 n_F(\omega_{e^-})[1-n_F(\omega_\nu)] \\
&= \frac{G_F^2}{2^5\pi^3}\frac{\left|\bar{u}^{(\omega_\nu)}_{\sigma_\nu}\gamma^0 u^{(\omega_{e^-})}_{\sigma_{e^-}}\right|^2\delta(\omega_{e^-}-\omega_\nu-\Delta m)}{e^{\pi\widetilde{\Delta m}}\cosh\tilde{\omega}_{e^-}\pi\cosh\tilde{\omega}_\nu\pi}, \qquad \text{(62a)}
\end{aligned}$$

$$\begin{aligned}
&\frac{1}{T}\frac{d^6\mathcal{P}^{p\to n}_{\text{(ii)}}}{d\omega_{\bar{\nu}}d\omega_{e^+}d^2k_{\bar{\nu}}d^2k_{e^+}} \\
&= \frac{1}{T}\sum_{\sigma_{e^+}}\sum_{\sigma_{\bar{\nu}}}\left|\mathcal{A}^{p\to n}_{\text{(ii)}}\right|^2 n_F(\omega_{\bar{\nu}})[1-n_F(\omega_{e^+})] \\
&= \frac{G_F^2}{2^5\pi^3}\frac{\left|\bar{u}^{(\omega_{e^+})}_{\sigma_{e^+}}\gamma^0 u^{(\omega_{\bar{\nu}})}_{\sigma_{\bar{\nu}}}\right|^2\delta(\omega_{\bar{\nu}}-\omega_{e^+}-\Delta m)}{e^{\pi\widetilde{\Delta m}}\cosh\tilde{\omega}_{\bar{\nu}}\pi\cosh\tilde{\omega}_{e^+}\pi}, \qquad \text{(62b)}
\end{aligned}$$

$$\begin{aligned}
&\frac{1}{T}\frac{d^6\mathcal{P}^{p\to n}_{\text{(iii)}}}{d\omega_{e^-}d\omega_{\bar{\nu}}d^2k_{e^-}d^2k_{\bar{\nu}}} \\
&= \frac{1}{T}\sum_{\sigma_{e^-}}\sum_{\sigma_{\bar{\nu}}}\left|\mathcal{A}^{p\to n}_{\text{(iii)}}\right|^2 n_F(\omega_{e^-})n_F(\omega_{\bar{\nu}}) \\
&= \frac{G_F^2}{2^5\pi^3}\frac{\left|\bar{u}^{(-\omega_{\bar{\nu}})}_{-\sigma_{\bar{\nu}}}\gamma^0 u^{(\omega_{e^-})}_{\sigma_{e^-}}\right|^2\delta(\omega_{e^-}+\omega_{\bar{\nu}}-\Delta m)}{e^{\pi\widetilde{\Delta m}}\cosh\tilde{\omega}_{e^-}\pi\cosh\tilde{\omega}_{\bar{\nu}}\pi}, \qquad \text{(62c)}
\end{aligned}$$

where

$$n_F(\omega) \equiv \frac{1}{1+e^{2\pi\tilde{\omega}}} \qquad (63)$$

is the fermionic thermal factor and $T = 2\pi\delta(0)$ is total proper time of the proton.

We can obtain the completeness relations by

$$\sum_\sigma u^{(\omega)}_\sigma(u,\mathbf{w})\bar{u}^{(\omega)}_\sigma(u,\mathbf{w}) = N^2\gamma^0\left[2\tilde{\ell}^2\left|K_{i\tilde{\omega}+\frac{1}{2}}(\tilde{\ell})\right|^2 + i\tilde{m}\tilde{\ell}\left\{(\gamma^3-\gamma^0)K^2_{i\tilde{\omega}+\frac{1}{2}}(\tilde{\ell})+(\gamma^3+\gamma^0)K^2_{i\tilde{\omega}-\frac{1}{2}}(\tilde{\ell})\right\}\right]. \qquad (64)$$

Using these relations, a direct calculation yields the spin sum for the process (i) as

$$\begin{aligned}
&\sum_{\sigma_{e^-}}\sum_{\sigma_\nu}\left|\bar{u}^{(\omega_\nu)}_{\sigma_\nu}\gamma^0 u^{(\omega_{e^-})}_{\sigma_{e^-}}\right|^2 \\
&= \frac{2^2}{\pi^4}\cosh\tilde{\omega}_{e^-}\pi\cosh\tilde{\omega}_\nu\pi\left[\tilde{\ell}_{e^-}\tilde{\ell}_\nu\left|K_{i\tilde{\omega}_{e^-}+\frac{1}{2}}(\tilde{\ell}_{e^-})K_{i\tilde{\omega}_\nu+\frac{1}{2}}(\tilde{\ell}_\nu)\right|^2 + \tilde{m}_e\tilde{m}_\nu\,\mathrm{Re}\left\{K^2_{i\tilde{\omega}_{e^-}+\frac{1}{2}}(\tilde{\ell}_{e^-})K^2_{i\tilde{\omega}_\nu-\frac{1}{2}}(\tilde{\ell}_\nu)\right\}\right]. \qquad (65)
\end{aligned}$$

We can perform the analogous calculation for the processes (ii) and (iii) .

According to Eq. (65), the differential transition rate of the process (i) is given by

$$\begin{aligned}
&\frac{1}{T}\frac{d^6\mathcal{P}^{p\to n}_{\text{(i)}}}{d\omega_{e^-}d\omega_\nu d^2k_{e^-}d^2k_\nu} \\
&= \frac{G_F^2}{2^3\pi^7e^{\pi\widetilde{\Delta m}}}\delta(\tilde{\omega}_{e^-}-\tilde{\omega}_\nu-\widetilde{\Delta m})\left[\tilde{\ell}_{e^-}\tilde{\ell}_\nu\left|K_{i\tilde{\omega}_{e^-}+\frac{1}{2}}(\tilde{\ell}_{e^-})K_{i\tilde{\omega}_\nu+\frac{1}{2}}(\tilde{\ell}_\nu)\right|^2 + \tilde{m}_e\tilde{m}_\nu\mathrm{Re}\left\{K^2_{i\tilde{\omega}_{e^-}+\frac{1}{2}}(\tilde{\ell}_{e^-})K^2_{i\tilde{\omega}_\nu-\frac{1}{2}}(\tilde{\ell}_\nu)\right\}\right]. \qquad (66)
\end{aligned}$$

By using them, the cross sections for each processes can be obtained as

$$\begin{aligned}
\Gamma^{p\to n}_{\text{(i)}} &= \frac{1}{T}\int d\mathcal{P}^{p\to n}_{\text{(i)}} \\
&= \frac{2G_F^2}{\pi^7e^{\pi\widetilde{\Delta m}}}\int_{\widetilde{\Delta m}}^\infty d\tilde{\omega}_{e^-}\left[\int_0^\infty d^2\widetilde{k}_{e^-}\tilde{\ell}_{e^-}\left|K_{i\tilde{\omega}_{e^-}+\frac{1}{2}}(\tilde{\ell}_{e^-})\right|^2\int_0^\infty d^2\widetilde{k}_\nu\tilde{\ell}_\nu\left|K_{i(\tilde{\omega}_{e^-}-\widetilde{\Delta m})+\frac{1}{2}}(\tilde{\ell}_\nu)\right|^2\right. \\
&\qquad\left.+\tilde{m}_e\tilde{m}_\nu\mathrm{Re}\left\{\int_0^\infty d^2\widetilde{k}_{e^-}K^2_{i\tilde{\omega}_{e^-}+\frac{1}{2}}(\tilde{\ell}_{e^-})\int_0^\infty d^2\widetilde{k}_\nu K^2_{i(\tilde{\omega}_{e^-}-\widetilde{\Delta m})-\frac{1}{2}}(\tilde{\ell}_\nu)\right\}\right], \qquad \text{(67a)}
\end{aligned}$$

$$\begin{aligned}
\Gamma^{p\to n}_{\text{(ii)}} &= \frac{2G_F^2}{\pi^7e^{\pi\widetilde{\Delta m}}}\int_0^\infty d\tilde{\omega}_{e^+}\left[\int_0^\infty d^2\widetilde{k}_{e^+}\tilde{\ell}_{e^+}\left|K_{i\tilde{\omega}_{e^+}+\frac{1}{2}}(\tilde{\ell}_{e^+})\right|^2\int_0^\infty d^2\widetilde{k}_{\bar{\nu}}\tilde{\ell}_{\bar{\nu}}\left|K_{i(\tilde{\omega}_{e^+}-\widetilde{\Delta m})+\frac{1}{2}}(\tilde{\ell}_{\bar{\nu}})\right|^2\right. \\
&\qquad\left.+\tilde{m}_e\tilde{m}_\nu\mathrm{Re}\left\{\int_0^\infty d^2\widetilde{k}_{e^+}K^2_{i\tilde{\omega}_{e^+}+\frac{1}{2}}(\tilde{\ell}_{e^+})\int_0^\infty d^2\widetilde{k}_{\bar{\nu}}K^2_{i(\tilde{\omega}_{e^+}-\widetilde{\Delta m})-\frac{1}{2}}(\tilde{\ell}_{\bar{\nu}})\right\}\right], \qquad \text{(67b)}
\end{aligned}$$

$$\Gamma^{p\to n}_{\rm (iii)} = \frac{2G_F^2}{\pi^7 e^{\pi\widetilde{\Delta m}}}\int_0^{\widetilde{\Delta m}} d\tilde{\omega}_{e^-}\left[\int_0^\infty d^2\widetilde{k}_{e^-}\tilde{\ell}_{e^-}\left|K_{i\tilde{\omega}_{e^-}+\frac{1}{2}}(\tilde{\ell}_{e^-})\right|^2\int_0^\infty d^2\widetilde{k}_{\bar{\nu}}\tilde{\ell}_{\bar{\nu}}\left|K_{i(\tilde{\omega}_{e^-}-\widetilde{\Delta m})+\frac{1}{2}}(\tilde{\ell}_{\bar{\nu}})\right|^2\right.$$
$$\left.+\tilde{m}_e\tilde{m}_\nu {\rm Re}\left\{\int_0^\infty d^2\widetilde{k}_{e^-}K^2_{i\tilde{\omega}_{e^-}+\frac{1}{2}}(\tilde{\ell}_{e^-})\int_0^\infty d^2\widetilde{k}_{\bar{\nu}}K^2_{i(\tilde{\omega}_{e^-}-\widetilde{\Delta m})-\frac{1}{2}}(\tilde{\ell}_{\bar{\nu}})\right\}\right], \quad (67c)$$

By summing them we can lump together the integration, and the total cross section becomes simply in the form

$$\Gamma^{p\to n}_{\rm acc} = \Gamma^{p\to n}_{\rm (i)} + \Gamma^{p\to n}_{\rm (ii)} + \Gamma^{p\to n}_{\rm (iii)}$$
$$= \frac{2G_F^2}{\pi^7 e^{\pi\widetilde{\Delta m}}}\int_{-\infty}^\infty d\tilde{\omega}\left[\int_0^\infty d^2\widetilde{k}_e\tilde{\ell}_e\left|K_{i\tilde{\omega}+\frac{1}{2}}(\tilde{\ell}_e)\right|^2\int_0^\infty d^2\widetilde{k}_\nu\ell_\nu\left|K_{i(\tilde{\omega}-\widetilde{\Delta m})+\frac{1}{2}}(\tilde{\ell}_\nu)\right|^2\right.$$
$$\left.+\tilde{m}_e\tilde{m}_\nu {\rm Re}\left\{\int_0^\infty d^2\widetilde{k}_e K^2_{i\tilde{\omega}+\frac{1}{2}}(\tilde{\ell}_e)\int_0^\infty d^2\widetilde{k}_\nu K^2_{i(\tilde{\omega}-\widetilde{\Delta m})-\frac{1}{2}}(\tilde{\ell}_\nu)\right\}\right]. (68)$$

It is hard to deal with these integral of modified Bessel's. Of course, we can use Eq. (28) again like the case of Section I. But we use more useful formula (see, for example, p219 of Ref. [13] or Ref. [14]).

$$x^\sigma K_\nu(x)K_\mu(x) = \frac{\sqrt{\pi}}{2}G^{40}_{24}\left(x^2\left|\begin{matrix}\frac{1}{2}\sigma, \frac{1}{2}\sigma+\frac{1}{2}\\ \frac{1}{2}(\nu+\mu+\sigma), \frac{1}{2}(\nu-\mu+\sigma), \frac{1}{2}(-\nu+\mu+\sigma), \frac{1}{2}(-\nu-\mu+\sigma)\end{matrix}\right.\right), \quad (69)$$

and by using the definition of G function, the integrand can be represented by power of ℓ and we can easily integrate with respect to ℓ.

We find

$$\Gamma^{p\to n}_{\rm acc} = \frac{G_F^2}{2^3\pi^4 e^{\pi\widetilde{\Delta m}}}\int_{C_s}\frac{ds}{2\pi i}\int_{C_t}\frac{dt}{2\pi i}\int_{-\infty}^\infty d\tilde{\omega}\frac{\tilde{m}_e^{2t+1}\tilde{m}_\nu^{2s+1}}{(2s+1)(2t+1)\Gamma(-s+1)\Gamma(-t+1)}$$
$$\times\left[\Gamma(-s+\frac{1}{2})\Gamma(-t+\frac{1}{2})\left|\Gamma(-s+i(\tilde{\omega}-\widetilde{\Delta m})+1)\Gamma(-t+i\tilde{\omega}+1)\right|^2\right.$$
$$\left.+\tilde{m}_e\tilde{m}_\nu\Gamma(-s-\frac{1}{2})\Gamma(-t-\frac{1}{2}){\rm Re}\left\{\Gamma(-s+i(\tilde{\omega}-\widetilde{\Delta m})+1)\Gamma(-s-i(\tilde{\omega}-\widetilde{\Delta m}))\Gamma(-t+i\tilde{\omega}+1)\Gamma(-t-i\tilde{\omega})\right\}\right], (70)$$

where all poles of complex s and t planes are picked up with C_s and C_t, respectively, by definition of G function.

To integrate with respect to ω, we use the formula of Barnes [14]:

$$\int_{-i\infty}^{i\infty} d\tilde{\omega}\ \Gamma(a+\tilde{\omega})\Gamma(b+\tilde{\omega})\Gamma(c-\tilde{\omega})\Gamma(d-\tilde{\omega}) = 2\pi i\frac{\Gamma(a+c)\Gamma(a+d)\Gamma(b+c)\Gamma(b+d)}{\Gamma(a+b+c+d)}$$
$$[{\rm Re}\ a, {\rm Re}\ b, {\rm Re}\ c, {\rm Re}\ d > 0]. \quad (71)$$

Eventually, we find that the total cross section in the Rindler frame is

$$\Gamma^{p\to n}_{\rm acc} = \frac{a^5 G_F^2}{2^5\pi^{\frac{7}{2}}e^{\pi\widetilde{\Delta m}}}\int_{C_s}\frac{ds}{2\pi i}\int_{C_t}\frac{dt}{2\pi i}\frac{(\tilde{m}_e^2)^s(\tilde{m}_\nu^2)^t}{\Gamma(-s-t+3)\Gamma(-s-t+\frac{7}{2})}$$
$$\times\left[\left|\Gamma(-s-t+i\widetilde{\Delta m}+3)\right|^2\Gamma(-s)\Gamma(-t)\Gamma(-s+2)\Gamma(-t+2)\right.$$
$$\left.+{\rm Re}\left\{\Gamma(-s-t+i\widetilde{\Delta m}+2)\Gamma(-s-t-i\widetilde{\Delta m}+4)\right\}\Gamma(-s+\frac{1}{2})\Gamma(-t+\frac{1}{2})\Gamma(-s+\frac{3}{2})\Gamma(-t+\frac{3}{2})\right]. \quad (72)$$

Comparing this to the results in inertial frame (31), we find that resulting expression agrees perfectly. This result shows the existence of Unruh effect is inevitable.

IV. DISCUSSIONS

We have analyzed the β decay and the inverse β decay of the accelerated proton in both frames. We found analytic expressions for both frame and found that they



FIG. 1: All residues of $\Gamma(-s)$ and $\Gamma(-s-\mu)$ are picked up.

agree with each other. If you see the calculation of G. E. A. Matsas and D. A. T. Vanzella in two-dimensional model [10, 11], you can realize that on four-dimensional model it became hopelessly complicated integral. So the main problem in this time is the complication of integral. To solve it, we used Barnes type representation as you can see from Eq. (28) and Eq. (69). We can demonstrate these formulae by picking up the poles in complex plane of the integral valuable. The sum of these readily becomes the infinity series which defines the special function. By using them, we accomplished perfectly analytical proof.

It is straightforward to apply our technique for two dimensional setup used in Ref. [10, 11], we can easily prove that the decay rates is independent of the frame.

APPENDIX A

In this appendix A we proof both key equations Eq. (28) and Eq. (30).

Firstly we derive Eq. (28). Through the path C_1, all poles of $\Gamma(-s)$ and $\Gamma(-s-\mu)$ are picked up (see FIG.1).

$$\begin{aligned}
&\frac{1}{2}\int_{C_1}\frac{ds}{2\pi i}\Gamma(-s)\Gamma(-s-\mu)\left(\frac{z}{2}\right)^{2s+\mu}\\
&=\frac{1}{2}\sum_{n=0}^{\infty}\frac{(-1)^n}{n!}\Gamma(\mp\mu-n)\left(\frac{z}{2}\right)^{2n\pm\mu}\\
&=\frac{1}{2}\sum_{n=0}^{\infty}\sum_{\pm}\frac{(-1)^n}{n!}\frac{\pi}{\Gamma(n\pm\mu+1)\sin(-n\mp\mu)\pi}\left(\frac{z}{2}\right)^{2n\pm\mu}\\
&=\frac{\pi}{2}\frac{-I_\mu(z)+I_{-\mu}(z)}{\sin\mu\pi}
\end{aligned}\tag{A1}$$

where $I_\mu(z)$ is modified Bessel function of the first kind. The last form is the definition of modified Bessel function $K_\mu(z)$ for non-integer μ. And you find the formula in case μ is integer n by setting $\mu=n$ after partial differentiation of this formula by μ.

Next we integrate Eq. (30) as FIG.2.



FIG. 2: The contour separates the poles of $\Gamma(-t)$ from those of $\Gamma(t-z)$.

$$\int_{C_2}\frac{dt}{2\pi i}\frac{\Gamma(-t)\Gamma(t-z)}{\Gamma(-z)}A^{-t+z}B^t$$

If you select the contour (i) you find

$$\begin{aligned}
&=A^z\sum_{n=0}^{\infty}\frac{(-1)^n\Gamma(-z+n)}{n!\Gamma(-z)}\left(\frac{B}{A}\right)^n\\
&=A^z\left(1+\frac{B}{A}\right)^z
\end{aligned}\tag{A2}$$

So this integration is the expansion form of $(A+B)^z$ in $B<A$. Similarly if you select the contour (ii) then you obtain the expansion form of $(A+B)^z$ in $B>A$. This integral representation of expansion of $(A+B)^z$ includes both cases of $B<A$ and $B>A$ by selecting the contour (i) and (ii), respectively.

APPENDIX B

In this appendix B, we show the explicit form of the integral. There are poles of the power of 1, 2 and 3.

The integral can be simply calculated by the change of variables to

$$s\to s-t,\quad t\to t.\tag{B1}$$

After this transformation we obtain

$$\begin{aligned}\Gamma^{p\to n} = \frac{a^5 G_F^2}{2^5 \pi^{\frac{7}{2}} e^{\pi\widetilde{\Delta m}}} \int_{C_t} \frac{dt}{2\pi i} \int_{C_s} \frac{ds}{2\pi i} \frac{(\tilde{m}_e^2)^{s-t} (\tilde{m}_\nu^2)^t}{\Gamma(-s+3)\Gamma(-s+\frac{7}{2})} \\ \times \Bigg[\Big| \Gamma(-s+i\widetilde{\Delta m}+3) \Big|^2 \Gamma(-t)\Gamma(-t+2)\Gamma(-s+t)\Gamma(-s+t+2) \\ + \mathrm{Re} \Big\{ \Gamma(-s+i\widetilde{\Delta m}+2)\Gamma(-s-i\widetilde{\Delta m}+4) \Big\} \Gamma(-t+\frac{1}{2})\Gamma(-t+\frac{3}{2})\Gamma(-s+t+\frac{1}{2})\Gamma(-s+t+\frac{3}{2}) \Bigg],\end{aligned} \tag{B2}$$

where $\psi(z) = \frac{d}{dz} \ln \Gamma(z)$ and C_t is the path separating the poles of, for example in first term, $\Gamma(-t)\Gamma(-t+2)$ from those of $\Gamma(-s+t)\Gamma(-s+t+2)$, and C_s is the path which picks up all the poles in s complex plane.

Firstly, we start by t integration because the existence of poles in t complex plane is independent of s. If we integrate by t, we obtain for $\tilde{m}_e > \tilde{m}_\nu$:

$$\begin{aligned}\Gamma^{p\to n} = \frac{a^5 G_F^2}{2^5 \pi^{\frac{7}{2}} e^{\pi\widetilde{\Delta m}}} \int \frac{ds}{2\pi i} \frac{\tilde{m}_e^{2s}}{\Gamma(-s+3)\Gamma(-s+\frac{7}{2})} \Bigg\{ \Bigg(\Gamma(-s)\Gamma(-s+2) + \left(\frac{\tilde{m}_\nu}{\tilde{m}_e}\right)^2 \Gamma(-s+1)\Gamma(-s+3) \\ + \sum_{n=2}^{\infty} \left(\frac{\tilde{m}_\nu}{\tilde{m}_e}\right)^{2n} \left[\psi(n-1) + \psi(n+1) - \psi(-s+n) - \psi(-s+n+2) - 2\ln\frac{\tilde{m}_\nu}{\tilde{m}_e} \right] \\ \times \frac{\Gamma(-s+n)\Gamma(-s+n+2)}{\Gamma(n-1)\Gamma(n+1)} \Bigg) \Big| \Gamma(-s+i\widetilde{\Delta m}+3) \Big|^2 \\ - \sum_{n=0}^{\infty} \left(\frac{\tilde{m}_\nu}{\tilde{m}_e}\right)^{2n+3} \left[\psi(n+1) + \psi(n+2) - \psi(-s+n+2) - \psi(-s+n+3) - 2\ln\frac{\tilde{m}_\nu}{\tilde{m}_e} \right] \\ \times \mathrm{Re} \Big\{ \Gamma(-s+i\widetilde{\Delta m}+2)\Gamma(-s-i\widetilde{\Delta m}+4) \Big\} \frac{\Gamma(-s+n+2)\Gamma(-s+n+3)}{\Gamma(n+1)\Gamma(n+2)} \Bigg\}\end{aligned} \tag{B3}$$

The simplest case is for massless neutrino. If we set $m_\nu = 0$, then only the first term remains non-vanishing. After a valuable transformation $s \to s + \frac{3}{2}$ we have

$$\begin{aligned}\Gamma^{p\to n}_{m_\nu=0} &= \frac{a^5 G_F^2 \tilde{m}_e^3}{2^5 \pi^{\frac{7}{2}} e^{\pi\widetilde{\Delta m}}} \int \frac{ds}{2\pi i} \frac{\Big| \Gamma(-s+i\widetilde{\Delta m}+\frac{3}{2}) \Big|^2 \Gamma(-s-\frac{3}{2})\Gamma(-s+\frac{1}{2})}{\Gamma(-s+\frac{3}{2})\Gamma(-s+2)} \tilde{m}_e^{2s} \\ &= \frac{G_F^2 m_e^3 a^2}{32\pi^{\frac{7}{2}} e^{\pi\widetilde{\Delta m}}} G^{40}_{24} \left(\frac{m_\nu^2}{a^2} \middle| \begin{matrix} \frac{3}{2}, \; 2 \\ \frac{1}{2}, \; -\frac{3}{2}, \; \frac{3}{2}+i\frac{\Delta m}{a}, \; \frac{3}{2}-i\frac{\Delta m}{a} \end{matrix} \right).\end{aligned} \tag{B4}$$

This is exactly the cross section obtained by Vanzella and Matsas with $c_V = 1$ and $c_A = 0$ (see Eq. (4.19) in [15]).

We perform the integration of Eq. (B3) with respect to s and find the following expansions of the cross section

$$
\begin{aligned}
\Gamma^{p\to n} &= \frac{a^5 G_F^2}{\pi^4 e^{\pi\widetilde{\Delta m}}} \\
&\times\Bigg\{\sum_{m=0}^{\infty}\sum_{\pm}\Bigg[\sum_{n=0}^{1}(-1)^{n+1}+\sum_{n=2}^{\infty}\frac{\psi(n-1)+\psi(n+1)-\psi(n-m\mp i\widetilde{\Delta m}-1)-\psi(n-m\mp i\widetilde{\Delta m}-3)-2\ln\frac{\tilde m_\nu}{\tilde m_e}}{\Gamma(n-1)\Gamma(n+1)}\Bigg] \\
&\qquad\qquad\times\frac{(-1)^m\Gamma(n-m\mp i\widetilde{\Delta m}-1)\Gamma(n-m\mp i\widetilde{\Delta m}-3)}{(-m\mp 2i\widetilde{\Delta m})B(m+1,-2m\mp 2i\widetilde{\Delta m})}\left(\frac{\tilde m_\nu}{\tilde m_e}\right)^{2n}\left(\frac{\tilde m_e}{2}\right)^{2(m\pm i\widetilde{\Delta m}+3)} \\
&-(1-\gamma)\left|\Gamma(i\widetilde{\Delta m}+1)\right|^2\left[\left(\frac{\tilde m_\nu}{2}\right)^4+\left(\frac{\tilde m_e}{2}\right)^4\right]+\sum_{k=3}^{\infty}\frac{\Gamma(2k-5)\left|\Gamma(-k+i\widetilde{\Delta m}+3)\right|^2}{\Gamma(k-1)\Gamma(k+1)}\left[\left(\frac{\tilde m_\nu}{2}\right)^{2k}+\left(\frac{\tilde m_e}{2}\right)^{2k}\right] \\
&-\sum_{k=2}^{\infty}\frac{\Gamma(2k-3)\left|\Gamma(-k+i\widetilde{\Delta m}+2)\right|^2}{\Gamma(k-1)\Gamma(k+1)}\left[\left(\frac{\tilde m_\nu}{2}\right)^{2k}\left(\frac{\tilde m_e}{2}\right)^{2}+\left(\frac{\tilde m_\nu}{2}\right)^{2}\left(\frac{\tilde m_e}{2}\right)^{2k}\right] \\
&+\sum_{n=2}^{\infty}\sum_{m=2}^{\infty}\left[-8\psi(2[n+m-3]+1)+2\sum_{\pm}\psi(-n-m\pm i\widetilde{\Delta m}+3)+\sum_{\pm}\psi(n\pm 1)+\sum_{\pm}\psi(m\pm 1)-2\ln\frac{\tilde m_\nu\tilde m_e}{4}\right] \\
&\qquad\qquad\times\frac{\Gamma(2[n+m-3]+1)\left|\Gamma(-n-m+i\widetilde{\Delta m}+3)\right|^2}{\Gamma(n-1)\Gamma(n+1)\Gamma(m-1)\Gamma(m+1)}\left(\frac{\tilde m_\nu}{2}\right)^{2n}\left(\frac{\tilde m_e}{2}\right)^{2m} \\
&+\sum_{n=0}^{\infty}\sum_{m=0}^{\infty}\Bigg(-\sum_{\pm}\left[\psi(n+1)+\psi(n+2)-\psi(n-m\mp i\widetilde{\Delta m})-\psi(n-m\mp i\widetilde{\Delta m}+1)-2\ln\frac{\tilde m_\nu}{\tilde m_e}\right] \\
&\qquad\qquad\times\frac{(-1)^m\Gamma(n-m\mp i\widetilde{\Delta m})\Gamma(n-m\mp i\widetilde{\Delta m}+1)}{2^2\Gamma(n+1)\Gamma(n+2)(-m\mp 2i\widetilde{\Delta m}+2)B(m+1,-2m\mp 2i\widetilde{\Delta m}+2)}\left(\frac{\tilde m_\nu}{\tilde m_e}\right)^{2n+3}\left(\frac{\tilde m_e}{2}\right)^{2(m\pm i\widetilde{\Delta m}+1)} \\
&-\sum_{\pm}\left[\psi(n+1)+\psi(n+2)-\psi(n-m\mp i\widetilde{\Delta m}-2)-\psi(n-m\mp i\widetilde{\Delta m}-1)-2\ln\frac{\tilde m_\nu}{\tilde m_e}\right] \\
&\qquad\qquad\times\frac{(-1)^m\Gamma(n-m\pm i\widetilde{\Delta m}-2)\Gamma(n-m\pm i\widetilde{\Delta m}-1)}{2^4\Gamma(n+1)\Gamma(n+2)(-m\pm 2i\widetilde{\Delta m}-2)B(m+1,-2m\pm 2i\widetilde{\Delta m}-2)}\left(\frac{\tilde m_\nu}{\tilde m_e}\right)^{2n+3}\left(\frac{\tilde m_e}{2}\right)^{2(m\mp i\widetilde{\Delta m}+4)} \\
&-\left[-8\psi(2[n+m]+1)+2\sum_{\pm}\psi(-n-m\pm i\widetilde{\Delta m}\mp 1)+\psi(n+1)+\psi(n+2)+\psi(m+1)+\psi(m+2)-2\ln\frac{\tilde m_\nu\tilde m_e}{4}\right] \\
&\qquad\qquad\qquad\times\frac{\Gamma(2[n+m]+1)\Gamma(-n-m+i\widetilde{\Delta m}-1)\Gamma(-n-m-i\widetilde{\Delta m}+1)}{\Gamma(n+1)\Gamma(n+2)\Gamma(m+1)\Gamma(m+2)}\left(\frac{\tilde m_\nu}{2}\right)^{2n+3}\left(\frac{\tilde m_e}{2}\right)^{2m+3}\Bigg)\Bigg\},
\end{aligned}
\tag{B5}
$$

where $B(p,q)=\frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$ and γ is Euler's constant.

This is the final form of the cross section. A natural question is that we have obtained the result which is not manifestly symmetric with respect to $\tilde m_\nu$ and $\tilde m_e$ although the original expression (31) is manifestly symmetric. The resolution of this puzzle is that the integral is of discontinuous type. Namely, we can obtain the result just by interchanging $\tilde m_\nu$ and $\tilde m_e$ for $\tilde m_e < \tilde m_\nu$. This can be checked directly by changing the order of integrations.

[1] S. W. Hawking, Nature (London) **248**, 30 (1974).

[2] W. G. Unruh, Phys. Rev. D **14**, 870 (1976).

[3] S. A. Fulling, Phys. Rev. D **7**, 2850 (1973).

[4] P. C. W. Davies, J. Phys. A: Gen. Phys. **8**, 609 (1975).

[5] G. L. Sewell, Ann. Phys. **141**, 201 (1982).

[6] N. D. Birrell and P. C. W. Davies, *Quantum Field Theory in Curved Space* (Cambridge University Press, Cambridge, England, 1982).

[7] P. Chen and T. Tajima, Phys. Rev. Lett. **83**, 256 (1999).

[8] R. Muller, Phys. Rev. D **56**, 953 (1997).

[9] C. Caso et al., The European Phys. Journ. C **3**, 1 (1998).

[10] G. E. A. Matsas and D. A. T. Vanzella, Phys. Rev. D **59**, 094004 (1999).

[11] D. A. T. Vanzella and G. E. A. Matsas, Phys. Rev. Lett. **87**, 151301 (2001).

[12] C. Itzykson and J.-B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).

[13] H. Bateman, *Higher Transcental Functions* (McGRAW-HILL, New York, 1953).

[14] I. S. Gradshteyn and I. M. Ryzhik, *Teble of Integrals, Series and Products* (Academic, New York, 1980).

[15] D. A. T. Vanzella and G. E. A. Matsas, Phys. Rev. D **63**, 014010 (2001).